

Polarized $\Lambda_b \rightarrow X_c \tau \nu$ in the SM and THDMChao-Shang Huang^{1,2} and Hua-Gang Yan¹¹ Institute of Theoretical Physics, Academia Sinica,
Beijing 100080, China² International Centre for Theoretical Physics,
P.O.Box 586, 34100 Trieste, Italy**Abstract**

The inclusive rate and τ spectrum for a polarized Λ_b -baryon to decay to charm hadronic final states and leptons $\tau \nu$ in the SM and a two-Higgs doublet model are computed. The $O(\alpha_s)$ QCD corrections to τ spectrum in the two-Higgs model are also given.

This work was supported in part by the National Natural Science Foundation of China and partly supported by Center of Chinese Advanced Science and Technology(CCAST).

I.Introduction

The semileptonic decay $B \rightarrow X_c l \nu$ has been extensively studied in both the standard model(SM) and a two-Higgs-doublet model(THDM) [1–9]. Compared with the B decay, in addition to the spectrum of the lepton arising from the decay, the various spin correlation quantities are of interest for the decay of a polarized Λ_b . It's well-known [10,11] that the heavy quarks produced in Z^0 decay are polarized and only charmed and beautiful Λ baryons seem to offer a practical method to measure the polarization of the corresponding heavy quark. The polarization transferred from a heavy quark Q to the corresponding Λ_Q is 100% [12] in the limit $m_Q \rightarrow \infty$. Thus the angular distributions of charged leptons [13,14] from semileptonic decays of Λ_b and Λ_c can be used as spin analysers for the decays of heavy quarks. Some aspects of the inclusive rate and l spectrum of polarized $\Lambda_b \rightarrow X_c l \nu$ have been studied in the SM [3,15,16].

The $\Lambda_b \rightarrow X_c \tau \nu$ (and $B \rightarrow X_c \tau \nu$) decay is sensitive to new physics, in particular, models with charged Higgs bosons. Because the charged Higgs bosons contribute at tree level, its contribution can not be cancelled by other new particles in the models. Therefore, the calculations of charged Higgs contributions with high accuracy will provide strong bound on parameters of the models when experimental measurement of the decay is available.

In this paper we investigate the polarized inclusive decay $\Lambda_b \rightarrow X_c \tau \nu$ in both the SM and THDM. In the SM we extend the results of Manohar and Wise [3] to the case of non-zero mass of the final state lepton (tau). We calculate the spin dependent form factors in the hadronic tensor to the $1/m_b^2$ order in HQET which do not contribute to the decay rate when the mass of the final state lepton is neglected. In the THDM we compute the inclusive rate and τ spectrum for a polarized $\Lambda \rightarrow X_c \tau \nu$ included the Λ_{QCD}^2/m_b^2 nonperturbative corrections and the $O(\alpha_s)$ perturbative corrections to τ spectrum.

The paper is organized as follows. In section II we calculate τ spectrum from a polarized Λ_b decay to the order $1/m_b^2$ in the $1/m_b$ expansion in the SM. Section III is devoted to calculations in the THDM. The $O(\alpha_s)$ QCD corrections to the decay are included. In section IV numerical results are given. Finally, a summary and discussions are presented in section V.

II. τ spectrum of $\Lambda_b \rightarrow X_c \tau \nu$ in the SM

We consider the inclusive semileptonic decay $\Lambda_b \rightarrow X_c \tau \nu$ in the SM. At the tree level, for unpolarized leptons the partial decay width can be written as

$$d\Gamma = \frac{G_f^2 |V_{cb}|^2}{(2\pi)^5 E_H} L^{\mu\nu} W_{\mu\nu} \frac{d^3 p_\tau}{2E_\tau} \frac{d^3 p_\nu}{2E_\nu}, \quad (1)$$

where E_H is the energy of Λ_b , $L_{\mu\nu}$ is the leptonic tensor

$$L^{\mu\nu} = 2(p_\tau^\mu p_\nu^\nu + p_\tau^\nu p_\nu^\mu - g^{\mu\nu} p_\tau \cdot p_\nu + i\epsilon^{\mu\nu\alpha\beta} (p_\tau)_\alpha (p_\nu)_\beta), \quad (2)$$

and $W_{\mu\nu}$ is the hadronic tensor

$$W_{\mu\nu} = (2\pi)^3 \sum_X \delta^4(p_{\Lambda_b} - q - p_X) \langle \Lambda_b(v, s) | J_\mu^\dagger | X \rangle \langle X | J_\nu | \Lambda_b(v, s) \rangle \quad (3)$$

with $p_{\Lambda_b} = m_{\Lambda_b} v$, $q = p_\tau + p_\nu$ and s being the spin of Λ_b . $J_\mu = \bar{c} \gamma_\mu \frac{(1-\gamma_5)}{2} b$ in eq.(3) is the hadronic current. The expansion of $W_{\mu\nu}$ in terms of Lorentz invariant structure functions W_i is defined by

$$W_{\mu\nu} = -g_{\mu\nu} W_1 + v_\mu v_\nu W_2 - i\epsilon_{\mu\nu\alpha\beta} v^\alpha q^\beta W_3 + q_\mu q_\nu W_4 + (q_\mu v_\nu + q_\nu v_\mu) W_5 + \{-q \cdot s [-g_{\mu\nu} W_1^s + v_\mu v_\nu W_2^s - i\epsilon_{\mu\nu\alpha\beta} v^\alpha q^\beta W_3^s + q_\mu q_\nu W_4^s + (q_\mu v_\nu + q_\nu v_\mu) W_5^s] + (s_\mu v_\nu + s_\nu v_\mu) W_6^s + (s_\mu q_\nu + s_\nu q_\mu) W_7^s + i\epsilon_{\mu\nu\alpha\beta} v^\alpha s^\beta W_8^s + i\epsilon_{\mu\nu\alpha\beta} q^\alpha s^\beta W_9^s\}. \quad (4)$$

The structure functions W_i can be calculated in heavy quark effective theory (HQET) [17] and results to the $1/m_b^2$ order are

$$W_1 = \delta(z) \left(\frac{m_b}{2} + \frac{K_b m_b}{6} - \frac{q \cdot v}{2} \right) - \delta'(z) \left(\frac{2 K_b m_b q^2}{3} + K_b m_b^2 q \cdot v - \frac{5 K_b m_b q \cdot v^2}{3} \right) + \delta''(z) \left(\frac{-2 K_b m_b^3 (q^2 - q \cdot v^2)}{3} + \frac{2 K_b m_b^2 q \cdot v (q^2 - q \cdot v^2)}{3} \right),$$

$$\begin{aligned}
W_2 &= \delta(z) \left(m_b + \frac{5 K_b m_b}{3} \right) - \frac{14 \delta'(z) K_b m_b^2 q.v}{3} + \delta''(z) \left(\frac{-4 K_b m_b^3 (q^2 - q.v^2)}{3} \right), \\
W_3 &= \frac{\delta(z)}{2} - \frac{5 \delta'(z) K_b m_b q.v}{3} + \delta''(z) \left(\frac{-2 K_b m_b^2 (q^2 - q.v^2)}{3} \right), \\
W_4 &= \frac{-4 \delta'(z) K_b m_b}{3}, \\
W_5 &= \frac{-\delta(z)}{2} - \delta'(z) \left(\frac{-4 K_b m_b^2}{3} - \frac{5 K_b m_b q.v}{3} \right) + \delta''(z) \left(\frac{2 K_b m_b^2 (q^2 - q.v^2)}{3} \right), \\
W_1^s &= \frac{-\delta(z) (1 + \epsilon_b)}{2} - \delta'(z) \frac{5 K_b m_b (-q.v)}{3} + \frac{2 \delta''(z) K_b m_b^2 (q^2 - q.v^2)}{3}, \\
W_2^s &= \frac{-4 \delta'(z) K_b m_b^2}{3}, \\
W_3^s &= \frac{-2 \delta'(z) K_b m_b}{3}, \\
W_4^s &= 0, \\
W_5^s &= \frac{2 \delta'(z) K_b m_b}{3}, \\
W_6^s &= \delta(z) \left(\frac{-(1 + \epsilon_b) m_b}{2} - \frac{5 K_b m_b}{6} \right) - \delta'(z) \left(-\frac{5 K_b m_b^2 q.v}{3} \right) + \frac{2 \delta''(z) K_b m_b^3 (q^2 - q.v^2)}{3}, \\
W_7^s &= \frac{\delta(z) (1 + \epsilon_b)}{2} - \delta'(z) \left(\frac{5 K_b m_b^2}{3} - K_b m_b (m_b - q.v) \right) - \\
&\quad \frac{2 \delta''(z) K_b m_b^2 (q^2 - q.v^2)}{3}, \\
W_8^s &= \delta(z) \left(\frac{(1 + \epsilon_b) m_b}{2} + \frac{K_b m_b}{6} \right) - \delta'(z) \left(\frac{5 K_b m_b^2 q.v}{3} \right) - \frac{2 \delta''(z) K_b m_b^3 (q^2 - q.v^2)}{3}, \\
W_9^s &= \frac{-\delta(z) (1 + \epsilon_b)}{2} - \delta'(z) \left(\frac{-5 K_b m_b^2}{3} + K_b m_b (m_b - q.v) \right) + \\
&\quad \frac{2 \delta''(z) K_b m_b^2 (q^2 - q.v^2)}{3}.
\end{aligned} \tag{5}$$

where

$$K_b = -\langle \Lambda_b(v, s) | \bar{b}_v \frac{(iD)^2}{2m_b^2} b_v | \Lambda_b(v, s) \rangle, z = (m_b v - q)^2 - m_c^2, \tag{6}$$

and ϵ_b is defined by [3]

$$\langle \Lambda_b(v, s) | \bar{b} \gamma^\lambda \gamma_5 b | \Lambda_b(v, s) \rangle = (1 + \epsilon_b) \bar{u}(v, s) \gamma^\lambda \gamma_5 u(v, s). \tag{7}$$

K_b and ϵ_b are only unknown two parameters at $O(m_b^{-2})$ for the Λ_b decay which parametrize the nonperturbative phenomena and are expected to be of order $(\Lambda_{QCD}/m_b)^2$. We will discuss them in section IV. The another parameter at $O(m_b^{-2})$

$$G_b = Z_b \langle H_b(v, s) | \bar{b}_v \frac{g G_{\alpha\beta} \sigma^{\alpha\beta}}{4m_b^2} b_v | H_b(v, s) \rangle \tag{8}$$

is equal to zero for $H_b = \Lambda_b$ due to the zero spin of the light degrees of freedom inside Λ_b . W_i (i=1,2,3) and W_i^s (i=1,2,3,6,8,9) have already been given by Manohar and Wise [3] and W_i (i=4,5) by Balk et al. [5]. We list them here only for completeness. From eqs(1),(2),(4) and(5) we get the differential decay rate

$$\frac{d\Gamma_W}{\Gamma_b dt dx dy d\cos\theta} = \hat{A}(x, t, y, \eta, \epsilon) + \hat{B}(x, t, y, \eta, \epsilon) \cos\theta, \tag{9}$$

here

$$\begin{aligned}
\hat{A}(x, t, y, \eta, \epsilon) = & [-12ty + 12xy - 12y\eta + K_b(-16t + 20xy + 16\eta)]\delta(z) - \\
& 4K_b(4t^2 - 4tx - 4ty - 5txy + 7x^2y - 5ty^2 + 7xy^2 + 4x\eta \\
& - 4y\eta - 5xy\eta - 5y^2\eta - 4\eta^2)\delta'(z) + 4K_by(4t - x^2 - 2xy \\
& - y^2)(t - x + \eta)\delta''(z), \\
\hat{B}(x, t, y, \eta, \epsilon) = & \frac{1}{\sqrt{x^2 - 4\eta}}\{[24t^2 - 24tx - 12txy + 12x^2y + 24x\eta - 12xy\eta - \\
& 24\eta^2 + K_b(-24tx + 20x^2y + 24x\eta - 32y\eta) + \epsilon_b(24t^2 - 24tx \\
& - 12txy + 12x^2y + 24x\eta - 12xy\eta - 24\eta^2)]\delta(z) - 4K_b(8t^2 + \\
& 6t^2x - 10tx^2 + 10t^2y - 18txy - 5tx^2y + 7x^3y - 5txy^2 + \\
& 7x^2y^2 + 10x^2\eta + 8ty\eta + 2xy\eta - 5x^2y\eta - 5xy^2\eta - 8\eta^2 - \\
& 6x\eta^2 - 2y\eta^2)\delta'(z) + 4K_b(4t - x^2 - 2xy - y^2)(t - x + \eta)(\\
& - 2t + xy + 2\eta)\delta''(z)\}
\end{aligned} \tag{10}$$

where

$$\Gamma_b = \frac{G_F^2 |V_{cb}|^2 m_b^5}{192\pi^3}, x = \frac{2E_\tau}{m_b}, y = \frac{2E_\nu}{m_b}, t = \frac{q^2}{m_b^2}, \eta = \frac{m_\tau^2}{m_b^2}, \epsilon = \frac{m_c}{m_b}$$

and θ is the angle between the τ direction and the Λ_b spin in the rest frame of the Λ_b . After integrating over t and y , one obtains the τ energy spectrum

$$\frac{d\Gamma_W}{\Gamma_b dx d\cos\theta} = A_W(x, \eta, \epsilon) + B_W(x, \eta, \epsilon) \cos\theta, \tag{11}$$

where $A_W(x, \eta, \epsilon)$ and $B_W(x, \eta, \epsilon)$ are given in Appendix. The $\eta \rightarrow 0$ limits of A_W and B_W agree with the results of Manohar and Wise [3]. The total inclusive decay width of $\Lambda_b \rightarrow X_c \tau \nu$ can be obtained by integrating the spectrum formula over range

$$2\sqrt{\eta} \leq x \leq 1 - \rho + \eta,$$

the result is not present here because one can easily obtain it from ref. [5] by taking $G_b=0$. Perturbative $O(\alpha_s)$ QCD corrections to the double differential distribution of the τ energy and the invariant mass of the lepton system for $b \rightarrow c \tau \nu$ have been studied by M. Jezabek et al. [18,19]. We will use eq.(30) in ref. [19] in our numerical analysis for the nonpolarized distribution of the τ energy.

III. τ spectrum of $\Lambda_b \rightarrow X_c \tau \nu$ in THDM

We consider the THDM [20–22] in which the up-type quarks get masses from Yukawa couplings to the one Higgs doublet H_2 (with the vacuum expectation value v_2) and down-type quarks and leptons get masses from Yukawa couplings to the another Higgs doublet H_1 (with the vacuum expectation value v_1). Such a model occurs as a natural feature in supersymmetric theories. For the sake of simplicity we shall use the Feynman rules of the THDM in MSSM [23]. In a THDM there are three diagrams contributing to the decay τ spectrum of $\Lambda_b \rightarrow X_c \tau \nu$ which correspond to W-exchange, Goldstone boson-exchange, and charged Higgs boson-exchange respectively if one uses a non-physical gauge. It is shown in ref. [8] that in the Landau gauge the rate can be decomposed into the sum of two incoherent decays:

$$M = M_W + M_S \Rightarrow |M|^2 = |M_W|^2 + |M_S|^2$$

where $M_S = M_G + M_H$. M_W , M_G and M_H are the W-mediated, Goldstone boson-mediated and Higgs boson-mediated decay amplitudes respectively. This decomposition has an advantage to simplify calculations, in particular, the calculations of QCD corrections. We assume the Landau gauge hereafter. Then the new thing needed to do is to calculate M_S .

For the purpose of calculating M_S the hadronic current J_μ in eq.(3) is replaced by

$$J_i = \bar{c}(a_i + b_i \gamma_5)b \quad (i = H, G) \tag{12}$$

with

$$a_H = m_b \tan \beta + m_c \cot \beta, b_H = m_b \tan \beta - m_c \cot \beta, a_G = -m_b + m_c, b_G = -m_b - m_c.$$

Following the same steps as those in section II, a straightforward calculation leads to

$$\frac{d\Gamma_H}{\Gamma_b dx d\cos\theta} = A_H(x, \eta, \epsilon, \xi, \tan\beta) + B_H(x, \eta, \epsilon, \xi, \tan\beta) \cos\theta, \quad (13)$$

$$\frac{d\Gamma_I}{\Gamma_b dx d\cos\theta} = A_I(x, \eta, \epsilon, \xi, \tan\beta) + B_I(x, \eta, \epsilon, \xi, \tan\beta) \cos\theta, \quad (14)$$

with $A_i(x, \eta, \epsilon, \xi, \tan\beta)$ and $B_i(x, \eta, \epsilon, \xi, \tan\beta)$ ($i = H, I$) given in Appendix. Here $d\Gamma_i/dx$ ($i = H, I$) denotes the contributions to the τ spectrum from Higgs-mediated and the interference term between Higgs-mediated and Goldstone boson-mediated respectively, and $\xi = m_H/m_b$.

For the spin independent terms A_H and A_I , our results agree with those obtained by Y. Grossman et al. [7]. Note that $B_I = 0$ at the leading order of $1/m_b$ expansion as $m_c \rightarrow 0$. This is due to the chiral difference of the vertices of W and Higgs. As we can see in numerical analysis that this makes B_I much smaller than B_W .

Combining eq.(11), eq.(13) and eq.(14), one obtains the τ spectrum of $\Lambda_b \rightarrow X_c \tau \nu$ in the THDM

$$\frac{d\Gamma_{THDM}}{\Gamma_b dx d\cos\theta} = (A_W + A_I + A_H) + (B_W + B_I + B_H) \cos\theta. \quad (15)$$

We now come to the position to calculate $O(\alpha_s)$ QCD corrections. Making use of the results obtained by A. Czarnecki et al. [24], Y. Grossman et al. got the $O(\alpha_s)$ corrections of the total width of $b \rightarrow c \tau \nu$ mediated by Higgs [8]. To get the $O(\alpha_s)$ corrections of τ spectrum one can also use their results. We find that the relation between $d\Gamma_{\alpha_s}^{H(I)}/dxdt$ and $d\Gamma_{\alpha_s}^{H(I)}/dt$ is very simple, as expected. $d\Gamma_{\alpha_s}^{H(I)}/dxdt$ is independent of x because Goldstone boson and Higgs are both scalar particles. Therefore, $d\Gamma_{\alpha_s}^{H(I)}/dxdt$ can be simply obtained by dividing $d\Gamma_{\alpha_s}^{H(I)}/dt$ by $x_{max} - x_{min}$, where x_{max}, x_{min} denote x 's kinematical upper and lower limits respectively. $d\Gamma_{\alpha_s}^{H(I)}/dt$ can be easily obtained from eq.(8) of ref. [24] by multiplying the lepton part. The results are

$$\begin{aligned} \frac{d\Gamma_{\alpha_s}^H}{dxdt} &= \frac{\sqrt{2}m_b^2\eta(t-\eta)\tan^2\beta}{16\pi^2\xi^4p_3}\Gamma(c_H^1, c_H^2, c_H^3), \\ \frac{d\Gamma_{\alpha_s}^I}{dxdt} &= \frac{\sqrt{2}m_b^2\eta(t-\eta)\tan\beta}{8\pi^2\xi^2tp_3}\Gamma(c_I^1, c_I^2, c_I^3), \end{aligned} \quad (16)$$

here

$$\begin{aligned} c_H^1 &= 2\tan^2\beta + 2\epsilon^2\cot^2\beta, c_H^2 = 4\epsilon, c_H^3 = \tan^2\beta - \epsilon^2\cot^2\beta, \\ c_I^1 &= -2\tan\beta + 2\epsilon^2\cot\beta, c_I^2 = 2\epsilon(\tan\beta - \cot\beta), c_I^3 = -\tan\beta - \epsilon^2\cot\beta, \end{aligned} \quad (17)$$

and $\Gamma(c^1, c^2, c^3)$ is [24]

$$\Gamma(c^1, c^2, c^3) = \frac{\alpha_s}{6\pi^2} \frac{G_F m_b^3 |V_{cb}|^2}{\sqrt{2}} [c^1 G_+ + c^2 G_- + c^3 G_0] \quad (18)$$

with

$$\begin{aligned} G_+ &= p_0 H + p_0 p_3 \left[\frac{9}{2} - 2 \ln\left(\frac{16p_3^4}{\epsilon^2 t}\right) \right] + \frac{1}{4t} Y_p (2 - t - 4t^2 + 3t^3 - 2\epsilon^2 - 2\epsilon^4 + 2\epsilon^6 - 4t\epsilon^2 - 5t\epsilon^4), \\ G_- &= H + p_3 \left[6 - 2 \ln\left(\frac{16p_3^4}{\epsilon^2 t}\right) \right] + \frac{1}{t} Y_p (1 - t - 2\epsilon^2 + \epsilon^4 - 3t\epsilon^2), \\ G_0 &= -6p_0 p_3 \ln \epsilon, \end{aligned}$$

where

$$H = 4p_0 [Li_2(p_+) - Li_2(p_-) - 2Li_2(1 - \frac{p_-}{p_+}) + \frac{1}{2} Y_p \ln(\frac{16p_3^4 t}{p_+^4}) - Y_w \ln \epsilon] + 2Y_w (1 - \epsilon^2) + \frac{2}{t} p_3 \ln \epsilon (1 + t - \epsilon^2),$$

$$p_0 \equiv \frac{1}{2}(1 - t + \epsilon^2), p_3 \equiv \frac{1}{2}\sqrt{1 + t^2 + \epsilon^4 - 2(t + \epsilon^2 + t\epsilon^2)}, p_{\pm} \equiv p_0 \pm p_3,$$

$$Y_p \equiv \frac{1}{2} \ln \frac{p_+}{P_-}, W_0 \equiv \frac{1}{2}(1+t-\epsilon^2), W_{\pm} \equiv W_0 \pm p_3, Y_w \equiv \frac{1}{2} \ln \frac{W_+}{W_-}.$$

After integrating (16) over t , we get τ spectrum numerically with only parameters $\tan\beta$ and ξ . As a check, we find that our numerical results of $\Gamma_{\alpha_s}^H$ and $\Gamma_{\alpha_s}^I$ agree with those obtained by Y. Grossman et al. [8].

IV. Numerical Results

In order to do numerical calculations we need to discuss values of parameters which are $K_b, \epsilon_b, m_c, m_b$ and the parameters in THDM.

1. Constraints on the parameters of THDM

In ref. [25,26], constraints on $\tan\beta$ from $K - \bar{K}$ and $B - \bar{B}$ mixing, $\Gamma(b \rightarrow s\gamma), \Gamma(b \rightarrow c\tau\nu)$ and R_b have been given

$$0.7 < \tan\beta < 0.6 \frac{m_{H_{\pm}}}{1\text{Gev}}$$

and also the lower limit $m_{H_{\pm}} > 200\text{Gev}$ has been given there. Taking the radiative correction and the $1/m_Q^2$ correction into account in the B meson decay, Y. Grossman et al. have a improved bound of R (defined by $R = \tan\beta/m_{H_{\pm}}$) which is

$$R < 0.49\text{Gev}^{-1}$$

. We will predict the τ spectrum and total width of Λ_b decay under these constraints.

2. About the parameters K_b, ϵ_b and m_b, m_c

K_b and ϵ_b characterize the $1/m_Q^2$ corrections to the decay distribution for $\Lambda_b \rightarrow X_c \tau \nu$ and are nonperturbative quantities independent of m_Q . Since quarks are not free physical particles, m_b and m_c can not be determined directly by experiment. However, we can estimate them by the phenomenological analysis of the heavy hadron spectra to the order $1/m_Q$. From the effective Lagrangian in HQET, the mass of a heavy hadron can be written as [30–32]

$$m_{h_Q} = m_Q + \bar{\Lambda}(j_l^P, I, S) + \frac{a(j_l^P, I, S)}{m_Q} + \frac{b(j_l^P, I, S)}{m_Q} \langle \vec{S}_Q \cdot \vec{j}_l \rangle + \dots, \quad (19)$$

where $\langle \vec{S}_Q \cdot \vec{j}_l \rangle = \frac{1}{2}[J(J+1) - j_l(j_l+1) - \frac{3}{4}]$ with J and j_l being the spins of the hadron and the light degrees of freedom inside the hadron respectively. The parameter $\bar{\Lambda}$ represents contributions come from the effective Lagrangian in the $m_Q \rightarrow \infty$ limit, and a, b are respectively associated with the kinetic energy and the color magnetic energy of the heavy quark inside the hadron. In present case, $a(0^+, 0, 0) = m_b^2 K_b$ and $b(0^+, 0, 0) = 0$. It is shown that $\epsilon_b \leq -2/3 K_b$ [29]. Furthermore, one can take $\epsilon_b = -2/3 K_b$ if one omits the contributions of terms arising from double insertions of chromomagnetic operator [27–29]. Starting from eq.(19), it is shown [32] that one can obtain $a(\frac{1}{2}^-, \frac{1}{2}, 0)$, m_c (and $\bar{\Lambda}(\frac{1}{2}^-, \frac{1}{2}, 0), b(\frac{1}{2}^-, \frac{1}{2}, 0)$) by using the observed masses of the doublets (B^*, B) and (D^*, D) if choosing m_b as input. Furthermore, $a(0^+, 0, 0)$ which is the parameter we need is determined by [32,3]

$$a(0^+, 0, 0) = \frac{h m_b}{1-h} [(m_{\Lambda_c} - \bar{m}_D) - (m_{\Lambda_b} - \bar{m}_B)] + a(\frac{1}{2}^-, \frac{1}{2}, 0), \quad (20)$$

where

$$h = \frac{m_{B^*} - m_B}{m_{D^*} - m_D}, \quad (21)$$

and $\bar{m}_H = \frac{1}{4}(m_H + 3m_{H^*})$, $H=B, D$. Therefore, if we choose $m_b = 5.1\text{Gev}$, the other parameters will be† :

$$m_c = h m_b = 1.65\text{Gev}$$

† the experimental data used in ref. [32] have been improved since then. Our data are from ref. [33].

$$K_b = \frac{0.142 m_c \text{Gev}}{m_b^2} \approx 0.009, \epsilon_b \approx -0.006$$

If we choose $m_b = 5.044$ which is a "critical" value based on eq.(19) [32], other parameters will be

$$m_c = 1.63 \text{Gev}, K_b \approx 0.006, \epsilon_b \approx -0.004$$

We will use these two sets of values and discriminate them by the first($m_b = 5.044 \text{Gev}$) and second($m_b = 5.1 \text{Gev}$) set respectively in the numerical computations.

Using the parameters given above, we obtain the total width in terms of $\tan\beta$ and m_H as follows:

$$\begin{aligned} \frac{\Gamma_W^1}{\Gamma_b} &= C_W^1 + D_W^1 \alpha_s, \\ \frac{\Gamma_H^1 + \Gamma_I^1}{\Gamma_b} &= C_H^1 + D_H^1 \alpha_s, \\ \frac{\Gamma_W^2}{\Gamma_b} &= C_W^2 + D_W^2 \alpha_s, \\ \frac{\Gamma_H^2 + \Gamma_I^2}{\Gamma_b} &= C_H^2 + D_H^2 \alpha_s, \end{aligned} \quad (22)$$

where

$$\begin{aligned} C_W^1 &= 0.109, \\ C_H^1 &= -\frac{0.0141 (0.253 + 1.16 \tan^2 \beta)}{\xi^2} + \frac{0.0141 (0.025 + 0.112 \tan^2 \beta + 0.239 \tan^4 \beta)}{\xi^4}, \\ D_W^1 &= -0.0476, \\ D_H^1 &= \frac{0.00804 (-0.165 - 0.577 \xi^2 - 0.374 \tan^2 \beta - \xi^2 \tan^2 \beta - 0.331 \tan^4 \beta)}{\xi^4}, \\ C_W^2 &= 0.112, \\ C_H^2 &= -\frac{0.0139 (0.262 + 1.19 \tan^2 \beta)}{\xi^2} + \frac{0.0139 (0.0256 + 0.107 \tan^2 \beta + 0.245 \tan^4 \beta)}{\xi^4}, \\ D_W^2 &= -0.0493, \\ D_H^2 &= \frac{0.0082 (-0.163 - 0.574 \xi^2 - 0.37 \tan^2 \beta - \xi^2 \tan^2 \beta - 0.328 \tan^4 \beta)}{\xi^4}, \end{aligned} \quad (23)$$

The superscript i (i=1, 2) in eqs.(22) and (23) denotes that the ith set of values of K_b, ϵ_b, m_c and m_b is used.

The explicit dependence of total width $\Gamma_{THDM} = \Gamma_W + \Gamma_H + \Gamma_I$ (normalized to the electron channel) on $\tan\beta$ and m_H is plotted in fig.1. It should be noted that the absolute value of Γ_{THDM} is very sensitive to m_b . Using the ratio between the width Γ_{THDM}^T and Γ_{THDM}^e separates the theoretical and experimental uncertainties and deletes the $|V_{cb}|^2 m_b^5$ factor which is not well-known yet. From fig.1 the following remarks can be drawn: (1) The normalized total width is not sensitive to the values of the parameters in HQET. The value of the normalized width for the second set of parameters is three percent larger than that for the first set. (2) In the range of $\tan\beta$ which is interesting physically, say, $\tan\beta < 60$, the normalized width changes roughly five to fifteen percent when $\tan\beta$ changes from twenty to sixty and $m_{H^\pm}$ is fixed. The normalized width changes the same order of magnitude for fixed $\tan\beta$ and changing $m_{H^\pm}$ from 200 to 400 GeV. From eqs. (22) and (23) it follows that the α_s corrections decrease the total width roughly 20 percent which is larger than that in SM. Because when $\tan\beta \gg 1$ C_H^i is proportional to $r^2 (-1 + 0.2 r^2)$ and D_H^i is proportional to $-r^2 (1 + 0.3 r^2)$ where $r = R m_b$, one can obtain constraints on r from the measurement of the total width.

τ spectrum for some typical values of $\tan\beta$ and m_H is calculated and the result is plotted in figs.2-5. The predictions in the SM are also plotted in the figs.. Here the α_s corrections for the spin-dependent term are not considered. We can see from figs.4-5 that the spin-dependent spectrum is quite different for $r \leq 1$ and $r \geq 2$ (Note that we have the constraints $r < 0.49 \text{Gev}^{-1} m_b \approx 2.5$ and $m_{H^\pm} \geq 200 \text{Gev}$ from experiments, as mentioned before). The reason is as follows. We know from Appendix that $B_H = -\frac{J r^2 \eta}{8 \xi^2} B_W \approx -r^4 \eta B_W / 4$. That is, it depends r^4 . For $r \leq 1$, $B_H \ll B_W$. As pointed in section III, B_I is negligibly small comparing with B_W due to the chiral difference of b quark couplings to W and $H_\pm$ which is deduced from Model II of

THDM. Therefore, the spectrum is almost the same as that in the SM, as can be seen from figs.4-5. For $r \geq 2$, B_H is as the same order of magnitude as B_W so that B_H and B_W tend to cancel each other, which makes the spin-dependent distribution of τ energy very small and a little dependent of the τ energy, as shown in figs.4-5. Thus one can say that if the τ spectrum is somewhat more isotropic than what the SM predicts, the THDM with large $\tan\beta$ (> 80) and $m_H \geq 200\text{Gev}$ is preferred in describing the nature. For the nonpolarized term, as can be seen from figs.4-5, the spectrum is very similar to that of B decay since the difference between the Λ_b decay and B decay comes from the $1/m_b^2$ corrections.

V. Summary

In summary, we have calculated the rate and τ spectrum of the inclusive semileptonic decay for a polarized $\Lambda_b \rightarrow X_c \tau \nu$ to the $1/m_b^2$ order in the $1/m_b$ expansion in the SM. The α_s corrections are included in the numerical computations for the spin independent terms of τ spectrum. Our results show that the spin dependent τ spectrum is significant enough to be seen.

We have also calculated the same quantities in a THDM. For the spin independent terms of τ spectrum arising from Higgs-mediated and the interference term, we have calculated the $O(\alpha_s)$ QCD corrections to the double differential distribution. Together with the α_s corrections in the SM given in ref. [18], we obtained all the α_s corrections to the non-polarized double differential distribution (and so the total width) in the THDM. The numerical results show that the branching ratio of $\Lambda_b \rightarrow X_c \tau \nu$ in THDM is of approximately 25 percent of that in the electron channel and the spin dependent τ spectrum can be used to estimate the size of $\tan\beta$ and $m_{H^\pm}$. The spectrum depends dominantly on R if $\tan\beta \gg 1$ so that from the measurement of the angular distribution of a polarized $\Lambda_b \rightarrow X_c \tau \nu$ in B-factories within the coming years one can obtain constraints on R.

It is obvious that substituting the u quark mass $m_u=0$ for m_c one immediately obtains the decay rate and τ spectrum for a polarized $\Lambda_b \rightarrow X_u \tau \nu$. And with minor changes one can extend the results in the paper to the inclusive semileptonic decay of a polarized $\Lambda_c \rightarrow X_{s,d} \tau \nu$. It is interesting to calculate the α_s corrections to the spin dependent term of τ spectrum.

Acknowledgments

One of the authors(H-G Yan) thanks Y. Grossman and F. Krüger for enlightening discussions. The other (C.-S. Huang) wishes to acknowledge International Centre for Theoretical Physics where the paper was written for the warm hospitality. We thank Y.P. Yao for kindly offering his unpublished article, L. Motyka and M. Jezabek for their program concerning the formulae of ref. [19].

-
- [1] J. Chay, H. Georgi and B. Grinstein, Phys.Lett.B247(1990)399.
 - [2] I.I. Bigi, M. Shifman, N.G. Uraltsev and A. Vainshtein, Phys.Rev.Lett.71(1993)496.
 - [3] A.V. Manohar and M.B. Wise, Phys.Rev.D49(1994)1310.
 - [4] A.F. Falk, Z. Ligeti, M. Neubert and Y. Nir, Phys.Lett.B326(1994)145.
 - [5] S. Balk, J.G. Körner, D. Pirjol and K. Schilcher, Z.Phys.C64(1994)37.
 - [6] B. Blok, L. Koyrakh, M. Shifman and A.I. Vainshtein, Phys.Rev.D49(1994)3356.
 - [7] Y. Grossman and Z. Ligeti, Phys.Lett.B332(1994)373.
 - [8] Y. Grossman, H.E. Haber and Y. Nir, Phys.Lett.B357(1995)630.
 - [9] M. Neubert, hep-ph/9702375, and references therein.
 - [10] J.H. Kühn and P.M. Zerwas, in *Heavy Flavours*, eds. A.J. Buras and M. Lindner, (World Scientific, Singapore, 1992), p.434
 - [11] J.H. Kühn et al., DESY Orange Report 92-123A(1992), Vol.I, p.255; J.H. Kühn, "Top Quark at a Linear Collider", in *Physics and Experiments with Linear e^+e^- Colliders*, eds.F.A. Harris et al., (World Scientific, Singapore, 1992), p.72.
 - [12] F.E. Close, J.G. Körner, R.J.N. Phillips and D.J. Summers, *J. Phys.*G18 (1992) 1716.
 - [13] G. Köpp, L.M. Sehgal and P.M. Zerwas, Nucl. Phys. B123(1977)77.
 - [14] B. Mele and G. Altarelli, Phys. Lett. B299(1993)345.
 - [15] M. Gremm, G. Köpp and L.M. Sehgal, Phys. Rev. D52 (1995) 1588.
 - [16] M. Gremm, G. Köpp, F. Krüger, and L.M. Sehgal, hep-ph/9510273, (published in the Proceedings of the EPS-HEP conference, Brussels, 1995).

- [17] N. Isgur and M. Wise, Phys.Lett.B232(1989)113; B237(1990)527; E. Eichten and B. Hill, Phys.Lett.B234(1990)511; H. Georgi, Phys.Lett.B240(1990)44; M. Neubert, Phys. Rep. 245(1994)259, and references therein.
- [18] M. Jeřabek, hep-hp/9410334 (Presented at XXVII International Conference on High Energy Physics 20-27, July 1994, Glasgow, Scotland), and references therein.
- [19] M. Jeřabek and L. Motyka, hep-ph/9701358.
- [20] L.J. Hall and M.B. Wise, Nucl. Phys. B187(1981)397; X.-G. He, T.D. Nguyen and R.R. Volkas, Phys. Rev D38(1988)814.
- [21] W. Skiba and J. Kalinowski, Nucl. Phys. B404(1983)3.
- [22] R.M. Barnett, G. Senjanovic and D. Wyler, Phys. Rev. D30(1984)1529.
- [23] J.F. Gunion, H.E. Haber, G. Kane and S. Dawson, The Higgs hunter's guide (Addison-Wesley, Reading, MA, 1990).
- [24] A. Czarnecki and S. Davidson, Phys. Rev. D48(1993)4183.
- [25] J. Kalinowski, Phys. Lett. B245(1990)201.
- [26] A.K. Grant, Phys. Rev. D51(1995)207.
- [27] A.F. Falk and M. Neubert, Phys. Rev. D47(1993)2982.
- [28] M. Gremm, F. Krüger and L.M. Sehgal, Phys. Lett. B355(1995)579.
- [29] J.G. Körner and D. Pirjol, Phys. Lett. B334(1994)399.
- [30] H.Y. Jin, C.S. Huang and Y.B. Dai, Z. Phys. C56(1992)702.
- [31] A.F. Falk and M. Neubert, Phys. Rev. D47(1993)2965, 2982.
- [32] C.S. Huang, Commun. Theor. Phys.24(1995)353.
- [33] Particle Data Group, Phys. Rev. D54(1996)1.

Appendix: Expressions of A_i and $B_i (i = W, H, I)$

$$\begin{aligned}
A_W = & \frac{\sqrt{x^2 - 4\eta}}{(1 - x + \eta)^3} (-1 + \epsilon^2 + x - \eta)^2 (3x + 3\epsilon^2 x - 5x^2 - \epsilon^2 x^2 + 2x^3 - \\
& 4\eta - 8\epsilon^2 \eta + 10x\eta + 3\epsilon^2 x\eta - 5x^2\eta - 4\eta^2 + 3x\eta^2) + \\
& 2K_b \frac{\sqrt{x^2 - 4\eta}}{3(1 - x + \eta)^5} (-5x^2 - 15\epsilon^4 x^2 + 20\epsilon^6 x^2 + 25x^3 + 21\epsilon^4 x^3 - \\
& 10\epsilon^6 x^3 - 50x^4 - 6\epsilon^4 x^4 + 2\epsilon^6 x^4 + 50x^5 - 25x^6 + 5x^7 + 14\eta + 6\epsilon^4 \eta \\
& - 20\epsilon^6 \eta - 70x\eta + 78\epsilon^4 x\eta - 80\epsilon^6 x\eta + 115x^2\eta - 147\epsilon^4 x^2\eta + \\
& 44\epsilon^6 x^2\eta - 40x^3\eta + 60\epsilon^4 x^3\eta - 10\epsilon^6 x^3\eta - 80x^4\eta - 6\epsilon^4 x^4\eta + 86x^5\eta \\
& - 25x^6\eta + 70\eta^2 - 126\epsilon^4 \eta^2 + 152\epsilon^6 \eta^2 - 280x\eta^2 + 300\epsilon^4 x\eta^2 - 80\epsilon^6 x\eta^2 \\
& + 370x^2\eta^2 - 147\epsilon^4 x^2\eta^2 + 20\epsilon^6 x^2\eta^2 - 130x^3\eta^2 + 21\epsilon^4 x^3\eta^2 - 80x^4\eta^2 \\
& + 50x^5\eta^2 + 140\eta^3 - 126\epsilon^4 \eta^3 - 20\epsilon^6 \eta^3 - 420x\eta^3 + 78\epsilon^4 x\eta^3 + 370x^2\eta^3 \\
& - 15\epsilon^4 x^2\eta^3 - 40x^3\eta^3 - 50x^4\eta^3 + 140\eta^4 + 6\epsilon^4 \eta^4 - 280x\eta^4 + 115x^2\eta^4 \\
& + 25x^3\eta^4 + 70\eta^5 - 70x\eta^5 - 5x^2\eta^5 + 14\eta^6), \tag{24}
\end{aligned}$$

$$\begin{aligned}
B_W = & \frac{(x^2 - 4\eta)}{(1 - x + \eta)^3} (-1 + \epsilon^2 + x - \eta)^2 (1 - \epsilon^2 - 3x - \epsilon^2 x + 2x^2 + 4\eta + \\
& 3\epsilon^2 \eta - 5x\eta + 3\eta^2) + \epsilon_b \frac{(x^2 - 4\eta)}{(1 - x + \eta)^3} (-1 + \epsilon^2 + x - \eta)^2 (1 - \epsilon^2 - 3x - \\
& \epsilon^2 x + 2x^2 + 4\eta + 3\epsilon^2 \eta - 5x\eta + 3\eta^2) + 2K_b \frac{(-x^2 + 4\eta)}{3(1 - x + \eta)^5} (5x - \\
& 15\epsilon^4 x + 10\epsilon^6 x - 25x^2 + 21\epsilon^4 x^2 + 4\epsilon^6 x^2 + 50x^3 - 6\epsilon^4 x^3 - 2\epsilon^6 x^3 \\
& - 50x^4 + 25x^5 - 5x^6 + 36\epsilon^4 \eta - 36\epsilon^6 \eta + 25x\eta - 51\epsilon^4 x\eta - 22\epsilon^6 x\eta \\
& - 100x^2\eta + 10\epsilon^6 x^2\eta + 150x^3\eta + 6\epsilon^4 x^3\eta - 100x^4\eta + 25x^5\eta + 60\epsilon^6 x\eta^2 \\
& + 50x\eta^2 + 51\epsilon^4 x\eta^2 - 20\epsilon^6 x\eta^2 - 150x^2\eta^2 - 21\epsilon^4 x^2\eta^2 + 150x^3\eta^2 \\
& - 50x^4\eta^2 - 36\epsilon^4 \eta^3 + 50x\eta^3 + 15\epsilon^4 x\eta^3 - 100x^2\eta^3 + 50x^3\eta^3 + 25x\eta^4 \\
& - 25x^2\eta^4 + 5x\eta^5). \tag{25}
\end{aligned}$$

$$A_H = \frac{\eta \tan^2 \beta \sqrt{x^2 - 4\eta}}{8\xi^4 (1 - x + \eta)^3} (1 - \epsilon^2 - x + \eta)^2 [L (6\epsilon x - 6\epsilon x^2 - 12\epsilon \eta +$$

$$\begin{aligned}
& 18\epsilon x\eta - 12\epsilon\eta^2) + F(3x + 3\epsilon^2x - 5x^2 - \epsilon^2x^2 + 2x^3 - 4\eta - \\
& 8\epsilon^2\eta + 10x\eta + 3\epsilon^2x\eta - 5x^2\eta - 4\eta^2 + 3x\eta^2)] + \\
& K_b\eta \frac{\tan^2\beta\sqrt{x^2-4\eta}}{12\xi^4(1-x+\eta)^5} [L(-18\epsilon x + 36\epsilon^3x - 18\epsilon^5x + 54\epsilon x^2 \\
& -120\epsilon^3x^2 + 66\epsilon^5x^2 - 36\epsilon x^3 + 132\epsilon^3x^3 - 60\epsilon^5x^3 - 36\epsilon x^4 \\
& -48\epsilon^3x^4 + 12\epsilon^5x^4 + 54\epsilon x^5 - 18\epsilon x^6 + 72\epsilon\eta - 96\epsilon^3\eta + 24\epsilon^5\eta \\
& -342\epsilon x\eta + 480\epsilon^3x\eta - 210\epsilon^5x\eta + 504\epsilon x^2\eta - 720\epsilon^3x^2\eta + \\
& 264\epsilon^5x^2\eta - 180\epsilon x^3\eta + 360\epsilon^3x^3\eta - 60\epsilon^5x^3\eta - 144\epsilon x^4\eta - \\
& 24\epsilon^3x^4\eta + 90\epsilon x^5\eta + 288\epsilon\eta^2 - 384\epsilon^3\eta^2 + 192\epsilon^5\eta^2 - 900\epsilon x\eta^2 \\
& 1080\epsilon^3x\eta^2 - 390\epsilon^5x\eta^2 + 756\epsilon x^2\eta^2 - 792\epsilon^3x^2\eta^2 + 102\epsilon^5x^2\eta^2 \\
& +36\epsilon x^3\eta^2 + 84\epsilon^3x^3\eta^2 - 180\epsilon x^4\eta^2 + 432\epsilon\eta^3 - 480\epsilon^3\eta^3 + 168\epsilon^5\eta^3 \\
& -828\epsilon x\eta^3 + 672\epsilon^3x\eta^3 - 54\epsilon^5x\eta^3 + 216\epsilon x^2\eta^3 - 96\epsilon^3x^2\eta^3 + \\
& 180\epsilon x^3\eta^3 + 288\epsilon\eta^4 - 192\epsilon^3\eta^4 - 234\epsilon x\eta^4 + 36\epsilon^3x\eta^4 - 90\epsilon x^2\eta^4 \\
& +72\epsilon\eta^5 + 18\epsilon x\eta^5) + F(-5x^2 - 15\epsilon^4x^2 + 20\epsilon^6x^2 + 25x^3 + 21\epsilon^4x^3 \\
& -10\epsilon^6x^3 - 50x^4 - 6\epsilon^4x^4 + 2\epsilon^6x^4 + 50x^5 - 25x^6 + 5x^7 + 14\eta + \\
& 6\epsilon^4\eta - 20\epsilon^6\eta - 70x\eta + 78\epsilon^4x\eta - 80\epsilon^6x\eta + 115x^2\eta - 147\epsilon^4x^2\eta \\
& +44\epsilon^6x^2\eta - 40x^3\eta + 60\epsilon^4x^3\eta - 10\epsilon^6x^3\eta - 80x^4\eta - 6\epsilon^4x^4\eta + \\
& 86x^5\eta - 25x^6\eta + 70\eta^2 - 126\epsilon^4\eta^2 + 152\epsilon^6\eta^2 - 280x\eta^2 + 300\epsilon^4x\eta^2 \\
& -80\epsilon^6x\eta^2 + 370x^2\eta^2 - 147\epsilon^4x^2\eta^2 + 20\epsilon^6x^2\eta^2 - 130x^3\eta^2 + 21\epsilon^4x^3\eta^2 \\
& -80x^4\eta^2 + 50x^5\eta^2 + 140\eta^3 - 126\epsilon^4\eta^3 - 20\epsilon^6\eta^3 - 420x\eta^3 + 78\epsilon^4x\eta^3 \\
& +370x^2\eta^3 - 15\epsilon^4x^2\eta^3 - 40x^3\eta^3 - 50x^4\eta^3 + 140\eta^4 + 6\epsilon^4\eta^4 - 280x\eta^4 \\
& +115x^2\eta^4 + 25x^3\eta^4 + 70\eta^5 - 70x\eta^5 - 5x^2\eta^5 + 14\eta^6)], \tag{26}
\end{aligned}$$

$$B_H = -\frac{Jr^2\eta}{8\xi^2} B_W, \tag{27}$$

$$\begin{aligned}
A_I = & -\frac{3\eta\sqrt{x^2-4\eta}(-1+\epsilon^2+x-\eta)^2}{\xi^2(1-x+\eta)^2} (2\epsilon^2 + 2\tan^2\beta - \epsilon^2x - 2x\tan^2\beta + \\
& 2\tan^2\beta\eta) - K_b\frac{\eta\sqrt{x^2-4\eta}}{\xi^2(1-x+\eta)^4} [6\epsilon^2 - 12\epsilon^4 + 6\epsilon^6 - 24\epsilon^2x + 24\epsilon^4x + \\
& 36\epsilon^2x^2 - 14\epsilon^4x^2 - 24\epsilon^2x^3 + 6\epsilon^2x^4 + 2\epsilon^4x^4 + 24\epsilon^2\eta - 4\epsilon^4\eta - 36\epsilon^6\eta \\
& -72\epsilon^2x\eta + 24\epsilon^6x\eta + 72\epsilon^2x^2\eta + 8\epsilon^4x^2\eta - 6\epsilon^6x^2\eta - 24\epsilon^2x^3\eta - 8\epsilon^4x^3\eta \\
& +36\epsilon^2\eta^2 - 4\epsilon^4\eta^2 + 6\epsilon^6\eta^2 - 72\epsilon^2x\eta^2 + 8\epsilon^4x\eta^2 + 36\epsilon^2x^2\eta^2 + 6\epsilon^4x^2\eta^2 \\
& +24\epsilon^2\eta^3 - 12\epsilon^4\eta^3 - 24\epsilon^2x\eta^3 + 6\epsilon^2\eta^4 + \tan^2\beta(-6 + 12\epsilon^2 - 6\epsilon^4 + 24x \\
& -48\epsilon^2x + 24\epsilon^4x - 36x^2 + 72\epsilon^2x^2 - 26\epsilon^4x^2 + 24x^3 - 48\epsilon^2x^3 + 8\epsilon^4x^3 \\
& -6x^4 + 12\epsilon^2x^4 - 30\eta + 48\epsilon^2\eta - 34\epsilon^4\eta + 96x\eta - 144\epsilon^2x\eta + 64\epsilon^4x\eta \\
& -108x^2\eta + 144\epsilon^2x^2\eta - 26\epsilon^4x^2\eta + 48x^3\eta - 48\epsilon^2x^3\eta - 6x^4\eta - 60\eta^2 \\
& +72\epsilon^2\eta^2 - 34\epsilon^4\eta^2 + 144x\eta^2 - 144\epsilon^2x\eta^2 + 24\epsilon^4x\eta^2 - 108x^2\eta^2 + \\
& 72\epsilon^2x^2\eta^2 + 24x^3\eta^2 - 60\eta^3 + 48\epsilon^2\eta^3 - 6\epsilon^4\eta^3 + 96x\eta^3 - 48\epsilon^2x\eta^3 - \\
& 36x^2\eta^3 - 30\eta^4 + 12\epsilon^2\eta^4 + 24x\eta^4 - 6\eta^5)], \tag{28}
\end{aligned}$$

$$\begin{aligned}
B_I = & \frac{3\eta\epsilon^2(-1+\epsilon^2+x-\eta)^2(x^2-4\eta)}{\xi^2(1-x+\eta)^2} + \epsilon_b\frac{3\eta\epsilon^2(-1+\epsilon^2+x-\eta)^2(x^2-4\eta)}{\xi^2(1-x+\eta)^2} \\
& - K_b\frac{\eta(1-\epsilon^2-x+\eta)}{\xi^2(1-x+\eta)^4} [-22\epsilon^2x + 20\epsilon^4x + 2\epsilon^6x + 62\epsilon^2x^2 - 30\epsilon^4x^2 - \\
& 8\epsilon^6x^2 - 52\epsilon^2x^3 + 4\epsilon^4x^3 + 4\epsilon^2x^4 + 6\epsilon^4x^4 + 10\epsilon^2x^5 - 2\epsilon^2x^6 + 92\epsilon^2\eta - \\
& 104\epsilon^4\eta + 12\epsilon^6\eta - 360\epsilon^2x\eta + 236\epsilon^4x\eta + 20\epsilon^6x\eta + 448\epsilon^2x^2\eta - 126\epsilon^4x^2\eta \\
& +8\epsilon^6x^2\eta - 176\epsilon^2x^3\eta - 4\epsilon^4x^3\eta - 2\epsilon^6x^3\eta - 12\epsilon^2x^4\eta + 2\epsilon^4x^4\eta + 8\epsilon^2x^5\eta \\
& +272\epsilon^2\eta^2 - 152\epsilon^4\eta^2 - 72\epsilon^6\eta^2 - 660\epsilon^2x\eta^2 + 156\epsilon^4x\eta^2 + 18\epsilon^6x\eta^2 + \\
& 420\epsilon^2x^2\eta^2 + 6\epsilon^4x^2\eta^2 - 4\epsilon^6x^2\eta^2 - 20\epsilon^2x^3\eta^2 - 8\epsilon^4x^3\eta^2 - 12\epsilon^2x^4\eta^2 + \\
& 264\epsilon^2\eta^3 - 56\epsilon^4\eta^3 + 12\epsilon^6\eta^3 - 328\epsilon^2x\eta^3 + 4\epsilon^4x\eta^3 + 32\epsilon^2x^2\eta^3 + 6\epsilon^4x^2\eta^3 \\
& +8\epsilon^2x^3\eta^3 + 80\epsilon^2\eta^4 - 8\epsilon^4\eta^4 - 6\epsilon^2x\eta^4 - 2\epsilon^2x^2\eta^4 - 4\epsilon^2\eta^5 + \tan^2\beta(2x - \\
& 4\epsilon^2x + 2\epsilon^4x - 6x^2 + 10\epsilon^2x^2 - 4\epsilon^4x^2 + 4x^3 - 6\epsilon^2x^3 + 2\epsilon^4x^3 + 4x^4 - \\
& 2\epsilon^2x^4 - 6x^5 + 2\epsilon^2x^5 + 2x^6 - 16\eta + 16\epsilon^2\eta + 74x\eta - 60\epsilon^2x\eta + 2\epsilon^4x\eta
\end{aligned}$$

$$\begin{aligned}
& -120 x^2 \eta + 66 \epsilon^2 x^2 \eta + 76 x^3 \eta - 16 \epsilon^2 x^3 \eta - 2 \epsilon^4 x^3 \eta - 8 x^4 \eta - 6 \epsilon^2 x^4 \eta - \\
& 6 x^5 \eta - 64 \eta^2 + 48 \epsilon^2 \eta^2 + 212 x \eta^2 - 108 \epsilon^2 x \eta^2 - 2 \epsilon^4 x \eta^2 - 228 x^2 \eta^2 + \\
& 54 \epsilon^2 x^2 \eta^2 + 4 \epsilon^4 x^2 \eta^2 + 76 x^3 \eta^2 + 6 \epsilon^2 x^3 \eta^2 + 4 x^4 \eta^2 - 96 \eta^3 + 48 \epsilon^2 \eta^3 + \\
& 212 x \eta^3 - 52 \epsilon^2 x \eta^3 - 2 \epsilon^4 x \eta^3 - 120 x^2 \eta^3 - 2 \epsilon^2 x^2 \eta^3 + 4 x^3 \eta^3 - 64 \eta^4 + \\
& 16 \epsilon^2 \eta^4 + 74 x \eta^4 - 6 x^2 \eta^4 - 16 \eta^5 + 2 x \eta^5 \Big],
\end{aligned} \tag{29}$$

where

$$F = 2 \tan^2 \beta + 2 \epsilon^2 \cot^2 \beta, J = 2 \tan^2 \beta - 2 \epsilon^2 \cot^2 \beta, L = 4 \epsilon.$$

When $\tan \beta \gg 1$, eqs. (3), (4) and (5) reduce to

$$A_H = \frac{r^4 \eta}{4} A_W + O(r^4 \tan \beta^{-2}), \tag{30}$$

$$B_H = -\frac{r^4 \eta}{4} B_W + O(r^4 \tan \beta^{-4}), \tag{31}$$

$$A_I = -\frac{6 \eta r^2 \sqrt{x^2 - 4 \eta}}{(1 - x + \eta)} (1 - x + \eta - \epsilon^2)^2 + O(r^2 \tan \beta^{-2}). \tag{32}$$


FIG. 1. Total width (normalized to the electron channel) in terms of $\tan \beta$ and m_H , using the (a) first ($m_b = 5.044 \text{ GeV}$), (b) second ($m_b = 5.1 \text{ GeV}$) set of value. The curves terminating at $\tan \beta = 100, 150, 200$ correspond to $m_H = 200 \text{ GeV}, 300 \text{ GeV}, 400 \text{ GeV}$ respectively.


FIG. 2. τ spectrum for different α_s and $\tan \beta$, $m_H = 200 \text{ GeV}$. The first set of parameter value ($m_b = 5.044 \text{ GeV}$) is used.


FIG. 3. τ spectrum for different α_s and $\tan \beta$, $m_H = 200 \text{ GeV}$. The second set of parameter value ($m_b = 5.1 \text{ GeV}$) is used.

FIG. 4. τ spectrum for different $\tan \beta$, $m_H = 200 \text{ GeV}$. The first set of parameter value ($m_b = 5.044 \text{ GeV}$) is used. $O(\alpha_s)$ corrections are not considered here.


FIG. 5. τ spectrum for different $\tan \beta$, $m_H = 200 \text{ GeV}$. The second set of parameter value ($m_b = 5.1 \text{ GeV}$) is used. $O(\alpha_s)$ corrections are not considered here.







(a)



(b)

